

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: A61K 31/59		A1	(11) International Publication Number: WO 96/31215
			(43) International Publication Date: 10 October 1996 (10.10.96)
(21) International Application Number: PCT/US96/04553		(81) Designated States: AU, BR, CA, CN, FI, HU, JP, KR, MX, NO, NZ, PL, SG, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 3 April 1996 (03.04.96)			
(30) Priority Data: 08/415,488 3 April 1995 (03.04.95) US		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(71) Applicant: BONE CARE INTERNATIONAL, INC. [US/US]; 313 West Beltline Highway, Madison, WI 53713 (US).			
(72) Inventors: KNUTSON, Joyce, C.; 24 North Prospect Avenue, Madison, WI 53705 (US). MAZESS, Richard, B.; 2526 Gregory Street, Madison, WI 53711 (US). BISHOP, Charles, W.; 5 LaPointe Terrace, Madison, WI 53719 (US).			
(74) Agents: WELCH, Teresa, J. et al.; Stroud, Stroud, Willink, Thompson & Howard, Suite 300, 25 West Main Street, P.O. Box 2236, Madison, WI 53701-2236 (US).			
(54) Title: USE OF VITAMIN D ₂ OR VITAMIN D ₄ -DERIVATIVES FOR THE MANUFACTURE OF A MEDICAMENT FOR THE TREATMENT OF SECONDARY HYPERPARATHYROIDISM			
(57) Abstract A method for preventing loss of bone mass or bone mineral content in a human being suffering from secondary hyperparathyroidism by administering a sufficient amount of 1 α -OH vitamin D ₂ , 1 α ,24(S)-(OH) ₂ vitamin D ₂ , 1 α -OH vitamin D ₄ or 1 α ,24(R)-(OH) ₂ vitamin D ₄ .			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

USE OF VITAMIN D2 OR VITAMIN D4-DERIVATIVES FOR THE MANUFACTURE OF A
MEDICAMENT FOR THE TREATMENT OF SECONDARY HYPERPARATHYROIDISM

5

This application is a continuation-in-part of U.S. Patent Application No. 08/119,895, which is a continuation of U.S. Patent Application No. 07/569,412, now U.S. Patent 5,104,864.

TECHNICAL FIELD

10

This invention relates generally to methods for treating and preventing metabolic bone disorders characterized by loss of bone mass or by disproportionately low bone mineral content. More specifically, this invention relates to a method for treating or preventing hyperparathyroidism secondary to end-stage renal disease, one of the concomitant results of which is the loss of bone mass or decreased mineral content (i.e., renal osteodystrophy).

15

BACKGROUND OF THE INVENTION

20

Numerous metabolic bone disorders are known which are characterized by loss of bone mass or bone mineral. These disorders include postmenopausal osteoporosis, senile osteoporosis, corticosteroid-induced osteopenia, anticonvulsant osteomalacia and renal osteodystrophy. Of these disorders, renal osteodystrophy is encountered in end-stage renal disease patients undergoing chronic dialysis.

25

As a group, these bone depletive disorders are a major and growing public health problem in the United States. Together, they cause more than 1 million bone fractures per year, primarily of the spine, hip, and distal forearm, and result in an annual cost above \$10 billion to the American society. Unfortunately, the incidence of these bone disorders will rise significantly as the mean age of the U.S. population continues to increase.

-2-

Despite differing etiologies, the aforementioned metabolic bone disorders develop during an extended period of negative calcium balance. This commonality of the disorders suggests that agents which stimulate intestinal calcium absorption and otherwise regulate calcium homeostasis may be effective in restoring calcium balance, thereby treating or preventing the development of such bone disorders.

It has long been known that vitamin D plays a critical role in stimulating calcium absorption and regulating calcium metabolism. The discovery of the active forms of vitamin D in the 1970's [M. F. Holick et al., *Proc. Natl. Acad. Sci. USA* 68, 803-804 (1971); G. Jones et al., *Biochemistry* 14, 1250-1256 (1975)] and active vitamin D analogues [M. F. Holick et al., *Science* 180, 190, 191 (1973); H. Y. Lam et al., *Science* 186, 1038-1040 (1974)], caused much excitement and speculation about the usefulness of these compounds in the treatment of bone depletive disorders.

Animal and early clinical studies examining the effects of these active vitamin D compounds suggested that such agents would be useful in restoring calcium balance. However, the best indicator of the efficacy of vitamin D compounds to prevent or treat depletive bone disorders is bone itself (or, in the case of renal osteodystrophy, serum levels of parathyroid hormone (PTH)) rather than calcium absorption or calcium balance. Certain clinical studies with $1\alpha,25-(\text{OH})_2$ vitamin D_3 , and $1\alpha\text{-OH}$ vitamin D_3 indicate that the ability of these agents to restore lost bone mass or bone mineral content is dose related. [See, S. M. Ott, C. H. Chesnut, *Annals of Int. Med.* 1989; 110:267-274; J. C. Gallagher et al., *Annals of Int. Med.* 1990; 113:649-655; J. Aloia et al., *Amer. J. Med.* 84:401-08 (1988)] M. Shiraki et al., *Endocrinol. Japan* 32, 305-315 (1985)].

These studies also indicate that at the dosage ranges required for these agents to be truly effective, toxicity in the form of hypercalcemia and hypercalciuria becomes a major problem. Attempts to increase the amount of $1\alpha,25-(\text{OH})_2$ vitamin D_3 above $0.5 \mu\text{g/day}$ have frequently resulted in toxicity. At dosage levels below $0.5 \mu\text{g/day}$, clinically significant effects are rarely

observed on bone. [See G. F. Jensen et al., *Clin. Endocrinol.* 16, 515-524 (1982); C. Christiansen et al., *Eur. J. Clin. Invest.* 11, 305-309 (1981)]. Doses of 2 $\mu\text{g/day}$ of $1\alpha\text{-OH}$ vitamin D_3 were found to have efficacy in increasing bone mass in patients exhibiting senile osteoporosis [O. H. Sorensen et al., *Clin. Endocrinol.* 7, 169S-175S (1977)]. Data from clinical studies in Japan, a population that has low calcium intake, indicate that efficacy is found with $1\alpha\text{-OH}$ vitamin D_3 when administered at 1 $\mu\text{g/day}$ [M. Shiraki et al., *Endocrinol. Japan.* 32:305-315 (1985); H. Orimo et al., *Bone and Mineral* 3, 47-52 (1987)]. However, at 2 $\mu\text{g/day}$, toxicity with $1\alpha\text{-OH}$ vitamin D_3 occurs in approximately 67 percent of the patients, and at 1 $\mu\text{g/day}$ this percentage is approximately 20 percent.

Thus, the prior art teaches that due to their toxicity, 1-hydroxylated vitamin D compounds can only be administered at dosages that are, at best, modestly beneficial in preventing or treating loss of bone or bone mineral content. Indeed, Aloia recommends that alternative routes of administration be sought which might avoid the toxicity problems and allow higher dosage levels to be achieved. [J. Aloia et al., *Amer. J. Med.* 84:401-408 (1988)].

Despite reported toxicities of $1\alpha\text{-OH}$ vitamin D_3 and $1\alpha,25\text{-(OH)}_2$ vitamin D_3 , these two compounds remain the drugs of choice for many bone depletive disease treatments. For example, in end stage renal disease, these two drugs remain the only approved forms of 1α -hydroxylated vitamin D for treating or preventing secondary hyperparathyroidism, although both drugs are not currently approved in all major pharmaceutical markets.

At present, in the United States, end stage renal disease afflicts approximately 200,000 individuals. In this disease, there is a progressive loss of cells of the proximal nephrons, the primary site for the synthesis of the vitamin D hormones (collectively " $1\alpha,25\text{-(OH)}_2\text{D}$ ") from 25-hydroxyvitamin D_3 and 25-hydroxyvitamin D_2 . In addition, the loss of functioning nephrons leads to retention of excess phosphorus which reduces the activity of the renal 25-hydroxyvitamin D- 1α -hydroxylase, the enzyme which catalyzes the reaction to produce the D hormones. These two events account for the low serum levels

of $1\alpha,25\text{-(OH)}_2\text{D}$ commonly found in patients with mild to moderate end stage renal disease.

5 Reduced serum levels of $1\alpha,25\text{-(OH)}_2\text{D}$ cause increased, and ultimately excessive, secretion of PTH by direct and indirect mechanisms. The resulting hyperparathyroidism leads to markedly increased bone turnover and its sequela of renal osteodystrophy, which may include a variety of other diseases, such as, osteitis fibrosa cystica, osteomalacia, osteoporosis, extraskeletal calcification and related disorders, e.g., bone pain, periarticular inflammation and Mockerberg's sclerosis. Reduced serum levels of $1\alpha,25\text{-(OH)}_2\text{D}$ also can cause muscle
10 weakness and growth retardation with skeletal deformities (most often seen in pediatric patients).

All previous clinical studies of hormonally active vitamin D drugs in end stage renal disease patients have focused exclusively on compounds derived from vitamin D_3 . Use of $1\alpha,25\text{-(OH)}_2\text{D}_3$ and $1\alpha\text{-OH-vitamin D}_3$ as replacement
15 therapy seeks to treat or prevent renal osteodystrophy by treating or preventing secondary hyperparathyroidism in end stage renal disease patients. As noted above, $1\alpha,25\text{-(OH)}_2\text{D}_3$ often causes toxic side effects (hypercalcemia and hyperphosphatemia) at dosages above $0.5\text{ }\mu\text{g}$, especially when concomitantly administered calcium phosphate binders are used to control serum phosphorus.
20 The minimum effective dose for preventing secondary hyperparathyroidism is in the range of 0.25 to $0.50\text{ }\mu\text{g/day}$; most patients respond to oral treatment doses of 0.5 to $1.0\text{ }\mu\text{g/day}$ or intravenous doses between 0.5 and $3.0\text{ }\mu\text{g}$ three times per week. As described above, the other commonly used vitamin D drug is $1\alpha\text{-OH-D}_3$, which often causes toxic effects at dosages over $1.0\text{ }\mu\text{g/day}$, especially when
25 used with calcium phosphate binders. The minimum effective dosage for preventing secondary hyperparathyroidism is in the range of 0.25 to $1.0\text{ }\mu\text{g/day}$, and most patients require treatment dosages of $1.0\text{ }\mu\text{g/day}$ or more. When either drug, $1\alpha,25\text{-(OH)}_2\text{D}_3$ or $1\alpha\text{-OH-D}_3$, is administered in higher dosages, both efficacy and toxicity are found to increase. Thus, the hormonally active
30 vitamin D_3 compounds are limited in their therapeutic usefulness due to their inherent toxicities.

-5-

To reduce the incidence of toxic side effects with $1\alpha,25-(\text{OH})_2\text{D}_3$ or $1\alpha\text{-OH-D}_3$, a low calcium dialysate with an ionized calcium concentration of 1.25 mM has been developed. However, it has been found that use of the low calcium dialysate has lead to higher serum PTH and phosphorus levels in patients who do not receive increased doses of oral calcium supplements and phosphate binders. When the dosages of calcium supplements and phosphate binders are increased, serum levels of phosphorus become controlled, but the incidence of hypercalcemia rises markedly. Thus, there are many problems associated with the use of current vitamin D therapies for secondary hyperparathyroidism.

Notwithstanding these known problems with use of the hormonally active vitamin D_3 for secondary hyperparathyroidism, the art has not adequately responded to date with the introduction of other vitamin compounds, derivatives or analogs that possess less inherent toxicity.

SUMMARY OF THE INVENTION

The present invention provides a method for treating or preventing hyperparathyroidism secondary to end stage renal disease by lowering (or maintaining low) serum parathyroid hormone levels in a patient suffering from the disease. The method at the same time ameliorates or prevents the renal osteodystrophy which can develop in such patients.

The foregoing, and other advantages of the present invention, are realized in one aspect thereof in a method for lowering serum (or plasma) PTH in patients suffering from hyperparathyroidism secondary to end stage renal disease, comprising: administering to these patients an effective amount of a vitamin D analog of formula (I), as described hereinbelow, to lower the serum PTH level.

The analog of formula (I) is any active vitamin D compound which has potent biological activity but low calcemic activity relative to the active forms of vitamin D_3 . Preferably such compounds are $1\alpha\text{-OH-vitamin D}_2$; $1\alpha,24(\text{S})\text{-(OH)}_2\text{-vitamin D}_2$; $1\alpha\text{-OH-vitamin D}_4$; or $1\alpha,24(\text{R})\text{-(OH)}_2\text{-vitamin D}_4$. The analog of formula (I) is administered in a dosage amount of 1 to about 100 $\mu\text{g/week}$. As used herein, the term "vitamin D analog" is meant to refer to compounds having

vitamin D hormonal bioactivity. It is also noted that a shorthand notation is often used for the D hormones, e.g., 1α -hydroxy vitamin D_2 may be referred to as 1α -OH-vitamin D_2 or simply 1α -OH- D_2 .

5 In another aspect, the invention is a pharmaceutical composition having serum (or plasma) PTH lowering activity, which includes, in unit dosage form, an effective amount of a vitamin D analog which is 1α -OH-vitamin D_2 , $1\alpha,24(S)$ -(OH) $_2$ -vitamin D_2 ; 1α -OH-vitamin D_4 ; or $1\alpha,24(R)$ -(OH) $_2$ -vitamin D_4 ; and a pharmaceutically acceptable excipient.

10 The treatment method of the present invention is an alternative to conventional therapy with $1\alpha,25$ -(OH) $_2$ vitamin D_3 or 1α -OH-vitamin D_3 ; the method is characterized by providing an active vitamin D compound having equivalent bioactivity but much lower toxicity than these conventional therapies. This is true especially in the case where oral calcium phosphate binders are used concomitantly to control serum phosphorus. As such, the method addresses a
15 long felt need in secondary hyperparathyroidism therapy.

A comparison of 1α -OH-vitamin D_2 to 1α -OH-vitamin D_3 has been conducted. 1α -OH-vitamin D_2 is equally active as 1α -OH-vitamin D_3 in the healing of rickets, in the stimulation of intestinal calcium absorption and in the elevation of serum inorganic phosphorous of rachitic rats. [G. Sjoden et al.,
20 *J. Nutr.* 114, 2043-2946 (1984)]. In the same laboratory animal, the inventors also have found that 1α -OH-vitamin D_2 is 5 to 15 times less toxic than 1α -OH-vitamin D_3 [see, also, G. Sjoden et al., *Proc. Soc. Exp. Biol. Med.* 178, 432-436 (1985)]. It has now been found that, for example, 1α -OH-vitamin D_2 may be safely administered for up to two years to human subjects experiencing
25 or having a tendency toward loss of bone mass or bone mineral content at dosages greater than 3 μ g/day.

The present invention is also intended to be used in all bone depletive disorders which respond to administration of active forms of vitamin D.

30 Other advantages and a fuller appreciation of specific adaptations, compositional variations, and physical attributes will be gained upon an examination of the following detailed description of preferred embodiments.

DETAILED DESCRIPTION


The present invention relates broadly to bone depletive disorders. However, the method of the present invention is most particularly adapted for use in ameliorating or preventing hyperparathyroidism secondary to end stage renal disease. The method also ameliorates or prevents the concomitant renal osteodystrophy of these patients with this disease. Accordingly, the present invention will now be described in detail with respect to such endeavors; however, those skilled in the art will appreciate that such a description of the invention is meant to be exemplary only and should not be viewed as limitative on the full scope thereof.

More specifically, the present invention relates to therapeutic methods for lowering the excessively high blood levels of parathyroid hormone (PTH) which are secondary to end stage renal disease. The method in accordance with the present invention has significantly less resultant hypercalcemia and hyperphosphatemia, especially in patients who use oral calcium phosphate binders to control serum phosphorus levels. These attributes are achieved through a novel treatment of a patient suffering from secondary hyperparathyroidism with a vitamin D analog of formula (I) as described hereinbelow.

In accordance with the invention, it has been found that when the analogs of formula (I) are administered to end stage renal disease patients with elevated serum parathyroid hormone, PTH concentration is lowered with significantly less hypercalcemia and hyperphosphatemia than is observed after the same amount of activated vitamin D administered in previously known formulations. Thus, the compounds of formula (I) have an improved therapeutic index relative to vitamin D₃ analogs.

-8-

The vitamin D analogs in accordance with the present invention have the general formula:



where B and C are each either H or together form a carbon-carbon double bond, and where R_1 is either a hydrogen or hydroxyl. The analogs of formula (I) are substantially less toxic than their vitamin D_3 counterparts when administered to patients experiencing hyperparathyroidism secondary to end stage renal disease. For patients using oral calcium phosphate binders, administration of the analogs of formula (I) at dosage levels higher than possible with the vitamin D_3 compounds provides greater efficacy than heretofore possible in treating secondary hyperparathyroidism.

Preferred among the analogs of formula (I) are: 1α -hydroxyvitamin D_2 (also known as 1α -hydroxyergocalciferol); 1α -hydroxyvitamin D_4 ; $1\alpha,24(S)$ -dihydroxyvitamin D_2 and $1\alpha,24(R)$ -dihydroxyvitamin D_4 . Most preferred is 1α -hydroxyvitamin D_2 , a prodrug for $1\alpha,25$ -dihydroxyvitamin D_2 which is an endogenous metabolite of vitamin D_2 .

It is noted that the medical community currently views vitamin D_3 compounds as biologically indistinguishable from the corresponding vitamin D_2 compounds. This is evident from the indiscriminate inclusion of either vitamin D_2 or D_3 in vitamin supplements prepared for human use, and from the interchangeable use of either vitamin in treating bone diseases caused by vitamin D deficiency. Curiously, medical experts consider the hormonally active

forms of the two vitamins to be equivalent despite lack of confirmation from a single human study. (It is also interestingly noted that vitamin D₄ is described in *The Merck Index* (Merck Index, 11th ed. (1989) p. 9932) as having doubtful biological activity.)

5 In parent application, Serial No. 08/119,895 and its parent application, now U.S. Patent 5,104,864, it has been shown that 1 α -OH-vitamin D₂ has the same biopotency as 1 α -OH-vitamin D₃ and 1 α ,25-(OH)₂-vitamin D₃ but is much less toxic. Even dosages up to 10 μ g/day of 1 α -OH-vitamin D₂ in women with postmenopausal osteoporosis (in both open label and double blind testing)
10 exhibited only mild hypercalciuria (> 300 mg/24 hrs), and marked hypercalcemia (> 11.0 mg/dL) solely due to 1 α -OH-vitamin D₂ was not evident. Additionally, the compound did not adversely affect kidney function, as determined by creatinine clearance and BUN; nor did it increase urinary excretion of hydroxyproline, indicating the absence of any stimulatory effect on bone
15 resorption. Administration of 1 α -OH-vitamin D₂ to healthy adult males in dosages up to 8 μ g/day showed no hypercalcemia or other adverse effects.

 The analogs of formula (I) are useful as active compounds in pharmaceutical compositions. The pharmacologically active analogs of this invention can be processed in accordance with conventional methods of pharmacy
20 to produce pharmaceutical agents for administration to patients, e.g., in admixtures with conventional excipients such as pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, enteral (e.g., oral) or topical application which do not deleteriously react with the active compounds. Suitable pharmaceutically acceptable carriers include but are not limited to water,
25 salt (buffer) solutions, alcohols, gum arabic, mineral and vegetable oils, benzyl alcohols, polyethylene glycols, gelatine, carbohydrates such as lactose, amylose or starch, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides, pentaerythritol fatty acid esters, hydroxy methylcellulose, polyvinyl pyrrolidone, etc. The pharmaceutical
30 preparations can be sterilized and if desired mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for

influencing osmotic pressure, buffers, coloring, flavoring and/or aromatic active compounds. If a solid carrier is used, the dosage form of the analogs may be tablets, capsules, powders, suppositories, or lozenges. If a liquid carrier is used, soft gelatin capsules, transdermal patches, aerosol sprays, topical creams, syrups
5 or liquid suspensions, emulsions or solutions may be the dosage form.

For parenteral application, particularly suitable are injectable, sterile solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants, including suppositories. Ampoules are convenient unit dosages. The dosage of the analogs in accordance with the present invention for parenteral
10 administration generally is about 1-30 μg given 1 to 3 times per week.

For enteral application, particularly suitable are tablets, dragées, liquids, drops, suppositories, or capsules. A syrup, elixir, or the like can be used wherein a sweetened vehicle is employed.

Sustained or directed release compositions can be formulated, e.g., liposomes or those wherein the active compound is protected with differentially
15 degradable coatings, e.g., by microencapsulation, multiple coatings, etc. It is also possible to freeze-dry the new compounds and use the lyophilizates obtained, for example, for the preparation of products for injection.

For topical application, there are employed as nonsprayable forms, viscous
20 to semi-solid or solid forms comprising a carrier compatible with topical application and having a dynamic viscosity preferably greater than water. Suitable formulations include but are not limited to solutions, suspensions, emulsions, creams, ointments, powders, liniments, salves, aerosols, etc., which are, if desired, sterilized or mixed with auxiliary agents, e.g., preservatives, etc.

Oral administration is preferred. Generally, the analogs of this invention
25 are dispensed by unit dosage form comprising about 0.25 to about 5.0 μg in a pharmaceutically acceptable carrier per unit dosage. The dosage of the analogs according to this invention generally is about 1 to about 100 $\mu\text{g}/\text{week}$, preferably about 3 to about 25 $\mu\text{g}/\text{week}$.

30 It will be appreciated that the actual preferred amounts of active analog in a specific case will vary according to the specific compound being utilized, the

-11-

particular compositions formulated, the mode of application, and the particular
situs being treated. Dosages can be determined using conventional
considerations, e.g., by customary comparison of the differential activities of the
subject compounds and of a known agent, e.g. by means of an appropriate
5 conventional pharmacological protocol.

The specific doses for each particular patient depend on a wide variety of
factors, for example, on the efficacy of the specific compound employed, on the
age, body weight, general state of health, sex, on the diet, on the timing and
mode of administration, on the rate of excretion, and on medicaments used in
10 combination and the severity of the particular disorder to which the therapy is
applied.

It is possible, if desired, to produce the metabolites of certain ones of the
analogs of formula (I), in particular by nonchemical means. For this purpose,
it is possible to convert them into a suitable form for administration together with
15 at least one vehicle or auxiliary and, where appropriate, combined with one or
more other active compounds.

The dosage forms may also contain adjuvants, such as preserving or
stabilizing adjuvants. They may also contain other therapeutically valuable
substances or may contain more than one of the compounds specified herein and
20 in the claims in admixture.

Bulk quantities of the vitamin D analogs for the practice of this invention
can be readily obtained in accordance with the processes of U.S. Patents
Nos. 3,907,843; 4,195,027; 4,202,829; 4,234,495; 4,260,549; 4,555,364; and
4,554,106 and U.S. Patent Application Serial Nos. 08/275,641 and 08/296,084.

25 As described hereinbefore, the analogs of formula (I) are preferably
administered to the human patients in oral dosage formulation. As an analog in
accordance with the present invention is released from the oral dosage
formulation, it is absorbed from the intestine into the blood.

The present invention is further explained by the following examples
30 which should not be construed by way of limiting the scope of the present
invention.

Example 1: Study Demonstrating Better Safety

The low toxicity of 1α -OH-vitamin D₂ in human patients was demonstrated in a clinical study involving 15 postmenopausal osteoporotic women. [*J. Bone Min. Res.*; 1994; 9:607-614.] The selected patients were
5 between 55 and 75 years of age, and exhibited L2-L3 vertebral bone mineral density ("BMD") between 0.7 and 1.05 g/cm², as determined by measurements with a LUNAR dual-photon absorptiometer. (The mean bone mineral density in women with osteoporosis is about 0.85 ± 0.17 g/cm², so that these limits correspond to about the 15th to 85th percentiles.)

10 On admission to the study, all patients received instruction on selecting a daily diet containing 400 to 600 mg of calcium. Compliance to this diet was verified at weekly intervals by 24-hour food records and by interviews with each patient.

15 All patients completed a one-week baseline period, a five- to seven-week treatment period, and a one-week post-treatment observation period. During the treatment period, patients orally self-administered 1α -OH-vitamin D₂ at an initial dose of 0.5 µg/day for the first week, and at successively higher doses of 1.0, 2.0, 4.0, 5.0, 8.0 and 10.0 µg/day in each of the following weeks. All doses were administered before breakfast.

20 Blood and urine chemistries were monitored on a weekly basis throughout the study. Key blood chemistries included fasting serum levels of calcium, phosphorus, osteocalcin, creatinine and blood urea nitrogen. Key urine chemistries included 24-hour excretion of calcium, phosphorus and creatinine.

25 Data from the study clearly demonstrated that 1α -OH-vitamin D₂ can be safely administered for short periods at high dose levels. In particular, the compound did not adversely affect kidney function, as determined by creatinine clearance and blood levels of urea nitrogen; nor did it increase urinary excretion of hydroxyproline, indicating the absence of any stimulatory effect on bone resorption. The compound had no effect on any routinely monitored serum
30 chemistries, indicating the absence of adverse metabolic effects.

-13-

A positive effect of 1α -OH-vitamin D₂ on calcium homeostasis was evident from dose-related increases observed in 24-hour urinary calcium levels, confirming that the compound increases intestinal calcium absorption, and from dose-related increases in serum osteocalcin, suggesting that the compound directly stimulates bone formation.

Example 2: Study Demonstrating Safety and Efficacy for Human Osteoporosis

The safety and efficacy of 1α -OH-vitamin D₂ as an oral treatment for osteoporosis was confirmed in a study involving 60 postmenopausal osteoporotic outpatients. The selected subjects had ages between 60 and 70 years, and exhibited L2-L3 vertebral BMD between 0.7 and 1.05 g/cm², as determined by dual-energy x-ray absorptiometry (DEXA). Exclusion criteria encompassed significant medical disorders and recent use of medications known to affect bone or calcium metabolism.

On admission to the study, each subject was assigned at random to one of two treatment groups; one group received up to a 104-week course of therapy with 1α -OH-vitamin D₂; the other received only placebo therapy. All subjects received instruction on selecting a daily diet containing 700-900 mg of calcium and were advised to adhere to this diet over the course of the study. Compliance to the diet was verified at regular intervals by 24-hour food records and by interviews with each subject.

During the treatment period, subjects from one group orally self-administered 1α -OH-vitamin D₂ at an initial dosage of 1.0 μ g/day for one week, and increased the dosage to 2.0, 3.0, 4.0 μ g/day in each of the following weeks, to a maximum dosage of 5.0 μ g/day. The dosage for any given subject was increased in this way until the rate of urinary calcium excretion was elevated to approximately 275-300 mg/24 hours, at which point the subject held the dosage constant at the highest level attained. Subjects from the second group self-administered a matching placebo medication every day, titrating the apparent

dosage upwards in the same manner as subjects being treated with 1α -OH-vitamin D₂.

5 Spinal and femoral neck BMD were measured in all subjects by DEXA at the beginning of the study, and at six-month intervals thereafter. Intestinal calcium absorption was estimated in all subjects by a single isotope technique at the beginning of the study, and at 12-month intervals. Serum levels of vitamin D metabolites were determined by radioreceptor binding assays at baseline and at six-month intervals. Serum osteocalcin, serum PTH and urine hydroxyproline also were determined at baseline and at six-month intervals.

10 Other blood and urine chemistries were monitored at regular intervals during the treatment period. These chemistries included serum calcium, serum ionized calcium, urine calcium, blood urea nitrogen, serum creatinine and creatinine clearance. Kidney-ureter-bladder (KUB) x-rays were obtained at baseline and at 12-month intervals thereafter.

15 The results of the study are summarized below:

Subjects: Sixty subjects enrolled in what was originally intended to be a 52-week study. Of these 60 subjects, 55 completed one year of treatment (28 active; 27 placebo); and 41 subjects completed an optional second year of treatment.

20 Test Drug Dosages: The average prescribed dosage for subjects who received 1α -OH-vitamin D₂ was 4.2 μ g/day at 52 weeks and 3.6 μ g/day at 104 weeks. The average prescribed dosage for placebo subjects was an apparent 4.8 μ g/day at 52 weeks and 4.8 μ g/day at 104 weeks.

25 Exclusions: One subject failed to comply with the prescribed dosage of test drug, as confirmed by an absence of serum $1\alpha,25$ -dihydroxyvitamin D₂ at any time during the study. Data for this subject were excluded from analysis. Three patients were diagnosed with hyperparathyroidism when the PTH assays were completed (in batch) at the study's conclusion; data for these subjects were excluded from analysis. No subjects were excluded from analysis for
30 noncompliance with the required dietary calcium intake of 700-900 mg/day.

Episodes of Hypercalcemia/Hypercalciuria: Marked hypercalcemia (> 10.8 mg/dL) occurred in one subject in association with an intercurrent illness. The prescribed dosage of 1α -OH-vitamin D₂ at the time of this episode was 5.0 µg/day. Moderate hypercalcemia (10.4-10.8 mg/dL) occurred in two subjects over the course of the study at prescribed dosages of 5.0 µg/day. Mild hypercalcemia (10.2-10.4 mg/dL) occurred in four subjects in the first year, and in two subjects in the second year. Hypercalciuria was observed occasionally over the two-year study in 17 subjects treated with 1α -OH-vitamin D₂.

Serum Calcium/Ionized Calcium: Mean serum calcium was approximately 0.1 to 0.2 mg/dL higher in subjects treated with 1α -OH-vitamin D₂ than in subjects treated with placebo. This difference was significant ($P < 0.05$) only during the second year of treatment. Mean serum ionized calcium was approximately 0.05 to 0.10 mg/dL higher in subjects treated with 1α -OH-vitamin D₂.

Urine Calcium: Mean urine calcium increased during the initial titration period in a dose-response fashion. After titration, mean urine calcium was 50 to 130% higher ($P < 0.001$) with 1α -OH-vitamin D₂ treatment than with placebo treatment.

Kidney Function: No significant changes were observed with long-term 1α -OH-vitamin D₂ treatment in BUN, serum creatinine or creatinine clearance. KUB x-rays revealed no abnormalities in either treatment group throughout the course of the study.

Bone: Bone mineral density (BMD) in the L2-L4 vertebrae progressively increased with 1α -OH-vitamin D₂ treatment and decreased with placebo treatment over the two-year study. The difference in spinal BMD between the treatment groups became statistically significant ($P < 0.05$) after 24 months of treatment. Similar changes were observed in femoral neck BMD with statistically significant differences observed after 18 months ($P < 0.001$) and 24 months ($P < 0.05$) of treatment.

Calcium Uptake: Intestinal absorption of orally administered ⁴⁵Ca increased by 40% ($P < 0.001$) after 52 weeks of 1α -OH-vitamin D₂ therapy, and

-16-

by 29% ($P < 0.5$) after 104 weeks of 1α -OH-vitamin D_2 therapy, relative to placebo control.

5 Vitamin D Metabolites: Treatment with 1α -OH-vitamin D_2 caused progressive increases in mean serum total $1\alpha,25$ -dihydroxyvitamin D from 21% ($P < 0.05$) at six months to 49% ($P < 0.01$) at 24 months relative to placebo therapy. This increase resulted from a dramatic rise in serum $1\alpha,25$ -dihydroxyvitamin D_2 which was partially offset by a 50+ % decrease in serum $1\alpha,25$ -dihydroxyvitamin D_3 . No treatment related changes were apparent in serum total 25-hydroxyvitamin D.

10 Biochemical Parameters:

Serum levels of PTH decreased with 1α -OH-vitamin D_2 therapy by 17% at 52 weeks and by 25% at 1-4 weeks, relative to placebo therapy.

15 Serum levels of osteocalcin were unchanged with long-term 1α -OH-vitamin D_2 therapy.

Fasting urine hydroxyproline:creatinine ratio tended to decrease with long-term 1α -OH-vitamin D_2 treatment but the observed differences between the 1α -OH-vitamin D_2 and placebo treatment groups were not significantly different.

20 The results of this study clearly indicated that 1α -OH-vitamin D_2 can be tolerated in higher long-term dosages than the commonly used vitamin D_3 analogues. They also showed that 1α -OH-vitamin D_2 is well tolerated in postmenopausal women at long-term dosages in the range of 2.0 to 3.0 $\mu\text{g/day}$, provided that individuals exhibiting abnormally high urine calcium levels (when
25 not receiving vitamin D therapy) are excluded from treatment. Long-term administration of such high dosages of 1α -OH-vitamin D_2 significantly reduced bone loss at the spine and femoral neck, the most frequent sites of osteoporotic fractures. These positive effects on bone were accompanied by a sustained increase in intestinal calcium absorption and a sustained decrease in serum PTH.
30 They were not accompanied by clear long-term trends in serum osteocalcin and urine hydroxyproline. Taken together, the results of this study demonstrate that

1α -OH-vitamin D₂ is safe and effective in the treatment of postmenopausal or senile osteoporosis.

Example 3: Open Label Study in End Stage Renal Disease Patients Exhibiting Secondary Hyperparathyroidism

5 Five end stage renal disease patients were enrolled in an open label study. The selected patients had ages between 36 and 72 years and had been on hemodialysis for at least 4 months prior to enrollment. The patients each had an average serum phosphorus in the range of 3.0 to less than or equal to 6.9 mg/dL during the two months prior to enrollment (often controlled by oral calcium phosphate binders), and had a history of elevated serum PTH values of greater
10 than 400 pg/mL when not receiving $1\alpha,25$ -(OH)₂-vitamin D₃ therapy.

Each patient had been receiving $1\alpha,25$ -(OH)₂ vitamin D₃ prior to enrollment, and discontinued the $1\alpha,25$ -(OH)₂ vitamin D₃ therapy for eight weeks prior to receiving 1α -OH-vitamin D₂. After 8 weeks, the patients received
15 treatment of 1α -OH-vitamin D₂ at a dosage of 4 µg/day for 6 weeks. Throughout the eight-week washout period and the treatment period, patients were monitored weekly or biweekly for serum intact PTH level and weekly for excessive elevation in serum levels of calcium and phosphorus.

Throughout the washout period and treatment period, patients underwent
20 routine hemodialysis (3 times per week) using a 1.25 mM calcium dialysate. They also ingested significant amounts of calcium phosphate binders (1-10g elemental Ca) to keep serum phosphorus levels below 6.9 mg/dL.

Baseline serum PTH was 480 ± 21 ; SCa (mg/dl), 9.8 ± 0.3 and serum phosphorus (mg/dl), 5.1 ± 0.2 . In three patients, serum PTH decreased by 68%,
25 74% and 87% after two weeks. In the other two patients, serum PTH declined by 33% in one and 3% in the other after four weeks. Overall, serum PTH decreased by $49 \pm 17\%$ and $33 \pm 9\%$ after two and four weeks of 1α -OH-vitamin D₂, respectively, ($p < 0.05$). Serum calcium was 10.2 ± 0.4 ($p < 0.05$) and 9.8 ± 0.2 (NS) and serum phosphorus was 5.4 ± 0.5 and 5.5 ± 0.8
30 at two and four weeks, respectively (NS). A rise in serum PTH from the second

to fourth weeks of 1α -OH-vitamin D_2 occurred when 1α -OH-vitamin D_2 was withheld in three patients with serum PTH < 130 ; they developed mild hypercalcemia (serum calcium, 10.3-11.4) that reversed after stopping 1α -OH-vitamin D_2 . No other adverse effects occurred. At 4-6 weeks of 1α -OH-vitamin D_2 treatment of 4 μ g, thrice weekly, four of five patients were in the target range of serum PTH; serum calcium was 10.0 ± 0.2 and serum phosphorus, 5.3 ± 0.2 mg/dl. The patient who failed to respond to six weeks of 1α -OH-vitamin D_2 treatment had a delayed response to both intravenous and oral calcitriol earlier, requiring several months of treatment before serum PTH fell. Serum PTH values in this patient fell by 38% after eight weeks of 1α -OH-vitamin D_2 treatment. These data show that 1α -OH-vitamin D_2 is efficacious and safe for the control of secondary hyperparathyroidism in end stage renal disease patients.

Example 4: Double Blind Study of Bone in End Stage Renal Disease Patients

A twelve-month double-blind placebo-controlled clinical trial is conducted with thirty-five men and women with renal disease who are undergoing chronic hemodialysis. All patients enter an eight-week control period during which time they receive a maintenance dose of vitamin D_3 (400 IU/day). After this control period, the patients are randomized into two treatment groups: one group receives a constant dosage of 1α -OH-vitamin D_2 (u.i.d.; a dosage greater than 3.0 μ g/day) and the other group receives a matching placebo. Both treatment groups receive a maintenance dosage of vitamin D_3 , maintain a normal intake of dietary calcium, and refrain from using calcium supplements. Oral calcium phosphate binders are used as necessary to maintain serum levels of phosphorus below 7.0 mg/dL. Efficacy is evaluated by pre- and post-treatment comparisons of the two patient groups with regard to (a) direct measurements of intestinal calcium absorption, (b) total body calcium retention, (c) radial and spinal bone mineral density, and (d) determinations of serum calcium and osteocalcin. Safety is evaluated by regular monitoring of serum calcium.

Analysis of the clinical data show that 1α -OH-vitamin D_2 significantly increases serum osteocalcin levels and intestinal calcium absorption, as determined by direct measurement using a double-isotope technique. Patients treated with this compound show normalized serum calcium levels, stable values for total body calcium, and stable radial and spinal bone densities relative to baseline values. In contrast, patients treated with placebo show frequent hypocalcemia, significant reductions in total body calcium and radial and spinal bone density. An insignificant incidence of hypercalcemia is observed in the treated group.

Example 5: Double-blind Study in End Stage Renal Disease (ESRD) Patients Exhibiting Secondary Hyperparathyroidism

Up to 120 ESRD (End Stage Renal Disease) patients undergoing chronic hemodialysis are studied in a multicenter, double-blind, placebo-controlled study based in two major U.S. metropolitan areas. The selected patients reside in two major metropolitan areas within the continental U.S., have ages between 20 and 75 years and have a history of secondary hyperparathyroidism. They have been on hemodialysis for at least four months, have a normal (or near normal) serum albumin, and have controlled serum phosphorus (often by using oral calcium phosphate binders).

On admission to the study, each patient is assigned at random to one of two treatment groups. One of these groups receives two consecutive 12-week courses of therapy with 1α -OH-vitamin D_2 ; the other receives a 12-week course of therapy with 1α -OH-vitamin D_2 followed, without interruption, by a 12-week course of placebo therapy. Each patient discontinues any $1\alpha,25$ -OH $_2$ -vitamin D_3 therapy for eight weeks prior to initiating 1α -OH-vitamin D_2 therapy ($4\mu\text{g/day}$). Throughout this eight-week washout (or control) period and the two subsequent 12-week treatment periods, patients are monitored weekly for serum calcium and phosphorus. Serum intact PTH is monitored weekly or biweekly, and bone-specific serum markers, serum vitamin D metabolites, serum albumin, blood chemistries, hemoglobin and hematocrit are monitored at selected intervals.

During the study, patients undergo routine hemodialysis (three times per week) using a 1.24 mM calcium dialysate and ingest calcium phosphate binders (such as calcium carbonate or calcium acetate) in an amount sufficient to keep serum phosphate controlled (≤ 6.9 mg/dL). Patients who develop persistent mild hypercalcemia or mild hyperphosphatemia during the treatment periods reduce their 1α -OH-vitamin D₂ dosage to 4 μ g three times per week (or lower). Patients who develop marked hypercalcemia or marked hyperphosphatemia immediately suspend treatment. Such patients are monitored at twice weekly intervals until the serum calcium or phosphorus is normalized, and resume 1α -OH-vitamin D₂ dosing at a rate which is 4 μ g three times per week (or lower).

During the eight-week washout period, the mean serum level of PTH increases progressively and significantly. After initiation of 1α -(OH)-vitamin D₂ dosing, mean serum PTH decreases significantly to less than 50% of pretreatment levels. Due to this drop in serum PTH, some patients need to reduce their dosage of 1α -OH-vitamin D₂ to 4 μ g three times per week (or to even lower levels) to prevent excessive suppression of serum PTH. In such patients, exhibiting excessive suppression of serum PTH, transient mild hypercalcemia is observed, which is corrected by appropriate reductions in 1α -OH-vitamin D₂ dosages.

At the end of the first 12-week treatment period, mean serum PTH is in the desired range of 130 to 240 pg/mL and serum levels of calcium and phosphorus are normal or near normal for end stage renal disease patients. At the end of the second 12-week treatment period (during which time 1α -OH-vitamin D₂ treatment is suspended and replaced by placebo therapy), mean serum PTH values markedly increase, reaching pretreatment levels. This study demonstrates that: (1) 1α -OH-vitamin D₂ is effective in reducing serum PTH levels, and (2) 1α -OH-vitamin D₂ is safer than currently used therapies, despite its higher dosages and concurrent use of high levels of oral calcium phosphate binder.

The foregoing examples demonstrate that 1α -OH-vitamin D₂ is effective in preventing or restoring the loss of bone mass or bone mineral content while

-21-

being substantially less toxic than $1\alpha,25-(\text{OH})_2$ -vitamin D_3 and 1α -OH-vitamin D_3 . It is to be understood that although the foregoing examples detail the use of 1α -OH-vitamin D_2 , other compounds within the scope of the claims may be readily utilized in the treatment of this invention with essentially equivalent results. For example, $1\alpha,24(\text{S})-(\text{OH})_2$ -vitamin D_2 shows activity equivalent to $1\alpha,24(\text{R})-(\text{OH})_2$ -vitamin D_2 and is also significantly less toxic than its vitamin D_3 counterpart. Also included within the scope of the claims would be administration of effective dosages of the analog of formula (I) in conjunction with administration of other hormones or other agents which have been shown to stimulate bone formation in subjects experiencing or tending toward loss of bone mass or bone mineral content.

Such hormones or other agents may include conjugated estrogens or their equivalents, calcitonin, biphosphonates, calcium supplements, cobalamin, pertussis toxin and boron. Possible dose ranges for these co-administered agents are provided in Table 1.

TABLE 1

Possible Oral Dose Ranges for Various Agents Co-Administered With 1α -Hydroxyvitamin D₂

5

	Agent	Dose Ranges		
		Broad	Preferred	Most Preferred
10	Conjugated Estrogens or Equivalent (mg/day)	0.3-5.0	0.4-2.4	0.6-1.2
	Sodium Fluoride (mg/day)	5-150	30-75	40-60
	Calcitonin (IU/day)	5-800	25-500	50-200
	Biphosphonates	50-2000	100-1500	250-1000
	Calcium Supplements (mg/day)	250-2500	500-1500	750-1000
15	Cobalamin (μ g/day)	5-200	20-100	30-50
	Pertussis Toxin (mg/day)	0.1-2000	10-1500	100-1000
	Boron (mg/day)	0.10-3000	1-250	2-100

20 Although the above examples detail dosage by mouth, it is to be understood that the compounds can also be administered in alternative fashions, including intranasally, transdermally, intrarectally, intravaginally, subcutaneously, intravenously, and intramuscularly.

25 In summary, the present invention provides therapeutic methods for lowering blood levels of parathyroid hormone which are secondary to end stage renal disease. The method in accordance with the present invention has significantly less resultant hypercalcemia and hyperphosphatemia.

30 While the present invention has now been described and exemplified with some specificity, those skilled in the art will appreciate the various modifications, including variations, additions, and omissions, that may be made in what has been described. Accordingly, it is intended that these modifications also be


-23-

encompassed by the present invention and that the scope of the present invention be limited solely by the broadest interpretation that lawfully can be accorded the appended claims.

-24-

CLAIMS:

1. A method for lowering or maintaining lowered serum parathyroid hormone in human patients suffering from hyperparathyroidism secondary to end stage renal disease, comprising: administering to said patients an effective amount of a vitamin D analog to lower and maintain lowered serum parathyroid hormone levels, said analog comprising formula (I):



wherein B and C are either hydrogen or a carbon-carbon double bond between C₂₂ and C₂₃; and R₁ is hydrogen or hydroxyl.

2. The method according to claim 1, wherein said analog of formula (I) is 1 α -OH-vitamin D₂; 1 α ,24(S)-(OH)₂-vitamin D₂; 1 α -OH-vitamin D₄; or 1 α ,24(R)-(OH)₂-vitamin D₄.
3. The method of claim 2 wherein said analog comprises a dosage of 1 to about 100 μ g/week.
4. The method of claim 1 wherein said analog, in solution, in a liquid vehicle ingestible by and nontoxic to said patients, is administered orally in encapsulated form.

-25-

5. The method of claim 1 wherein said analog is administered in combination with at least one agent characterized by said agent's ability to reduce loss of bone mass, or bone mineral content in patients.

5 6. The method of claim 5 wherein said agent includes other vitamin D compounds, conjugated estrogens, sodium fluorides, biphosphonates, cobalamin, pertussin toxin or boron.

7. The method of claim 1, wherein said administration of said analog is parenteral.

10 8. The method of claim 7 wherein said administration is by subcutaneous, intramuscular, or intravenous injection, nasopharyngeal or mucosal absorption, or transdermal absorption.

9. The method of claim 1 wherein said administration of said analog is nonparenteral.

15 10. A pharmaceutical composition having serum parathyroid hormone lowering activity, comprising, in unit dosage form, an effective amount of a vitamin D analog which is 1α -OH-vitamin D_2 ; $1\alpha,24(S)$ -(OH) $_2$ -vitamin D_2 ; 1α -OH-vitamin D_4 ; or $1\alpha,24(R)$ -(OH) $_2$ -vitamin D_4 ; and a pharmaceutically acceptable excipient.

20 11. The composition of claim 10, wherein said amount is 0.25 to 5.0 μ g.

12. A pharmaceutical composition as claimed in claim 10 which further comprises, in combination, at least one agent characterized by said agent's ability to reduce loss of bone mass or bone mineral content in mammals experiencing or tending toward said loss of bone mass or bone mineral content.

13. A pharmaceutical composition as claimed in claim 12 wherein said agent includes other vitamin D compounds, conjugated estrogens, calcitonin, sodium fluoride, bisphosphonates, calcium supplements, cobalamin, pertussin toxin or boron.

5 14. As an article of manufacture, a tablet having activity to lower parathyroid hormone level as measured by blood serum level of parathyroid hormone over time after ingestion, comprising: from about 0.25 to 5.0 μ g of a vitamin D analog selected from the group consisting of 1α -OH-vitamin D_2 ; $1\alpha,24(S)$ -(OH) $_2$ -vitamin D_2 ; 1α -OH-vitamin D_4 ; and $1\alpha,24(R)$ -(OH) $_2$ -vitamin D_4 .

10 15. A method for achieving an effect in a patient comprising administering an effective amount of the composition of claim 10 to the patient wherein the effect is lowering or maintaining lowered serum parathyroid hormone levels, and thus decreasing loss of bone mass or bone mineral content.

15 16. A method of treating a human to alleviate or prevent the pathological effects of hyperparathyroidism secondary to end stage renal disease, wherein the method comprises administering to said human a vitamin D analog selected from the group consisting of 1α -OH-vitamin D_2 ; $1\alpha,24(S)$ -(OH) $_2$ -vitamin D_2 ; 1α -OH-vitamin D_4 ; and $1\alpha,24(R)$ -(OH) $_2$ -vitamin D_4 wherein said compound is administered to said human in an amount sufficient to lower or
20 maintain lowered serum parathyroid hormone levels in said human to thereby alleviate or prevent said effects.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/04553

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/59

According to International Patent Classification (IPC) or to both national classification and IPC:

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	BIOCHEM. J., vol. 310, no. 1, 15 August 1995, pages 233-241, XP002010503 STRUGNELL S. ET AL.: "1alpha,24(S)-dihydroxyvitamin D2: a biologically active product of 1alpha-hydroxyvitamin D2 made in the human hepatoma, Hep3B" see abstract see page 233, left-hand column, line 14 - line 20 see page 234, left-hand column, last paragraph see page 240, right-hand column, last paragraph --- -/--	1-16

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

8 August 1996

Date of mailing of the international search report

29. 08. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (- 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (- 31-70) 340-3016

Authorized officer

Economou, D

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 96/04553

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	J.BONE MINER.RES., vol. 9, no. 5, May 1994, pages 607-614, XP000578685 GALLAGHER J.C. ET AL.: "Effects of increasing doses of lalpha-hydroxyvitamin D2 on calcium homeostasis in postmenopausal women" cited in the application see abstract see page 607, left-hand column, line 1 - page 608, left-hand column, line 27 see page 608, right-hand column, paragraph 3 see page 611, left-hand column, line 8 - line 9 see page 612, left-hand column, line 19 - line 21 see page 613, left-hand column, line 36 - line 39 ---	1-16
P,X	ENDOCRINOLOGY, vol. 136, no. 11, November 1995, pages 4749-4753, XP000575597 KNUTSON J.C. ET AL.: "Metabolism of lalpha-hydroxyvitamin D2 to activated dihydroxyvitamin D2 metabolites decreases endogenous lalpha,25-dihydroxyvitamin D3 in rats and monkeys" see abstract see page 4749, left-hand column, line 16 - right-hand column, line 3 see page 4750, left-hand column, line 5 - line 8 see page 4750, left-hand column, line 20 - line 23 ---	1-16
X	WO,A,92 05130 (LUNAR CO.) 2 April 1992 see page 1, paragraph 1 - paragraph 2 see page 4, paragraph 3 see page 5, last paragraph - page 8, line 32 see claims 13,14,27,28,30-33 ---	10-14
X	WO,A,92 12165 (LUNAR CO.) 23 July 1992 see page 1, paragraph 1 see page 7, line 1 - page 9, line 27 see claims 5-7,10,11,29 ---	10-14
X	WO,A,93 14763 (LUNAR CO.) 5 August 1993 see page 1, paragraph 1 see page 5, last paragraph - page 6, paragraph 1 see page 12 - page 13, paragraph 1; example 2 ---	10,11
	--- -/--	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/04553

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,94 16711 (LUNAR CO) 4 August 1994 see page 2, line 22 - page 3, line 15 see page 6, line 21 - line 30 ---	10-14
X	US,A,4 833 125 (NEER ET AL.) 23 May 1989 see column 3, line 55 - line 68 see column 6; table 1 see column 7, line 16 - column 8, line 12 see claims 1-8 ---	10-14
X	US,A,4 698 328 (NEER ET AL.) 6 October 1987 see column 3, line 55 - line 67 see column 5, line 18 - line 58 see column 6; table 1 see column 7, line 1 - line 43 ---	10-14
X	EP,A,0 503 630 (KURARAY CO., LTD.) 16 September 1992 see page 2, line 55 - page 3, line 1 ---	1,2
X	EP,A,0 562 497 (NISSHIN FLOUR MILLING CO. LTD.) 29 September 1993 see page 2, line 51 - page 3, line 15 see page 7, line 48 - page 8, line 27 see page 11; example 2 see claims 1,3-5 -----	10-14

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/04553

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9205130	02-04-92	AU-B- 650286	16-06-94
		AU-B- 8542291	15-04-92
		CA-A- 2069084	22-03-92
		CN-A- 1061220	20-05-92
		EP-A- 0503035	16-09-92
		NZ-A- 239897	26-03-96
		US-A- 5488120	30-01-96
WO-A-9212165	23-07-92	AU-B- 1247592	17-08-92
		CN-A- 1067243	23-12-92
		EP-A- 0550702	14-07-93
WO-A-9314763	05-08-93	AU-B- 3656193	01-09-93
		CA-A- 2129120	05-08-93
		EP-A- 0631500	04-01-95
		JP-T- 7503714	20-04-95
WO-A-9416711	04-08-94	US-A- 5350745	27-09-94
		CA-A- 2155009	04-08-94
		EP-A- 0680329	08-11-95
US-A-4833125	23-05-89	US-A- 4698328	06-10-87
US-A-4698328	06-10-87	US-A- 4833125	23-05-89
		AU-B- 599905	02-08-90
		CA-A- 1288695	10-09-91
		DE-A- 3686343	17-09-92
		EP-A- 0197514	15-10-86
		IE-B- 59620	09-03-94
		JP-B- 7072138	02-08-95
		JP-A- 62000033	06-01-87
		JP-A- 7179358	18-07-95
EP-A-0503630	16-09-92	IL-A- 101222	31-03-96
		JP-A- 6025039	01-02-94
		US-A- 5334740	02-08-94
EP-A-0562497	29-09-93	JP-A- 5320127	03-12-93